



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to 16.2 per thousand of the population. This was also higher than the rate for the same week of last year, in which it amounted to only 15.6. Of the large German towns and cities, however, two-thirds showed less favorable health conditions than Berlin, the following places having a considerably higher rate of mortality than this city, viz, Munich, Nuremberg, Stuttgart, Frankfort-on-the-Main, Bremen, Brunswick, Danzig, Königsberg, Breslau, Dresden, Cologne, Aix la Chapelle, as well as London, Paris, and Vienna. The death rate of Hamburg and Halle corresponded exactly with that of Berlin, while the following places had a lower rate of mortality than this city, viz, Charlottenburg (with 15.5); Schöneberg (with the minimum of 7 per thousand); Leipzig, Hanover, Kiel, Carlsruhe. Since the foregoing week there has been a considerable increase in the number of deaths among children in the first year of life. The infant mortality rate, 4.2 per year and thousand, was higher than the Hamburg and Leipzig figure, but lower than the Munich rate.

There were registered 111 deaths from phthisis pulmonalis; 73 deaths from acute diseases of the respiratory organs; 37 deaths from cancer; 11 deaths from scarlet fever; 9 deaths from diphtheria; 5 deaths from measles. Finally, 9 persons died by violence.

INDIA.

Report from Calcutta—Cholera and plague mortality—Plague increasing in Rangoon.

Acting Assistant Surgeon Eakins reports, March 23, as follows:

There were no transactions at this port during the week ended March 18, 1905.

During the week ended March 18, 1905, there were 30 deaths from cholera and 405 deaths from plague in Calcutta.

In Bengal, during the week ended March 11, 1905, there were 9,406 cases and 8,243 deaths from plague.

In India, during the week ended March 11, 1905, there were 52,504 cases and 45,541 deaths.

Plague increasing in Rangoon.

Plague is steadily increasing in Rangoon. The last information available, up to the 20th instant, reports 170 cases with 136 deaths.

A European photographer had plague, and the report, from a lay correspondent, states that he took the disease by photographing a plague patient, but is on a fair way to recovery.

JAPAN.

Report from Yokohama—Inspection of vessels—Communicable diseases—Emigrants recommended for rejection.

Passed Assistant Surgeon Moore reports, March 16, as follows:

During the week ended March 11, 1905, bills of health were issued to 2 steamships having an aggregate personnel of 159 crew and 164 passengers; 74 steerage passengers were bathed and their clothing was disinfected; 80 intending emigrants were examined with reference to